Molecular Brain Research, 3 (1988) 298–300
Elsevier

Author Index

Allen, J.M., Martin, J.B. and Heinrich, G., Neuropeptide Y gene expression in PC12 cells and its regulation by nerve growth factor: a model for developmental regulation, 39

Banner, C., Hwang, J.-J., Shapiro, R.A., Wenthold, R.J., Nakatani, Y., Lampel, K.A., Thomas, J.W., Huie, D. and Curthoys, N.P., Isolation of a cDNA for rat brain glutaminase, 247
Banner, C., see Kim, Y.S., 77
Bannon, M.J., Elliott, P.J. and Bunney, E.B., Striatal tachykinin biosynthesis: regulation of mRNA and peptide levels by dopamine agonists and antagonists, 31
Bech-Hansen, N.T., see McLachlan, D.R.C., 255
Bergeron, C., see McLachlan, D.R.C., 255
Biswal, N., Patel, A.G. and Max, S.R., Regulaton of viral and cellular genes in a human neuroblastoma cell line latently infected with herpes simplex virus type 2, 95
Black, I.B., see Friedman, W.J., 203
Blakely, R.D., see Guarda, A.S., 223
Bowcock, A.M., Fahnestock, M., Goslin, K. and Shooter, E.M., The NGF and kallikrein genes of mouse, the African rat *Mastomys natalensis* and man: their distribution and mode of expression in the salivary gland, 165
Brown, I.R., see Sprang, G.K., 89
Brugge, J.S., see Walaas, S.I., 215
Bunney, E.B., see Bannon, M.J., 31
Busciglio, J., see Cáceres, A., 233

Cáceres, A., Busciglio, J., Ferreira, A. and Steward, O., An immunocytochemical and biochemical study of the microtubule-associated protein MAP-2 during post-lesion dendritic remodeling in the central nervous system of adult rats, 233
Calvetta, L., see Sivam, S.P., 155
Carbini, L., see McCaman, R.E., 107
Civelli, O., see Sivam, S.P., 155
Conlon, J.M., see McGregor, G.P., 207
Cooper, B.J., see Roch, J.-M., 61
Coyle, J.T., see Guarda, A.S., 223
Curthoys, N.P., see Banner, C., 247

Danysz, W., see Kaczmarek, L., 183
Davidson, N., see Sutton, F., 187
Davis, L.G., see Manning, R.W., 293
Deguchi, T., see Saito, M., 83
Douglass, J., see Sivam, S.P., 155
Dreyfus, C.F., see Friedman, W.J., 203

Ebendal, T., see Läkfors, L., 55
Elliott, P.J., see Bannon, M.J., 31

Fahnestock, M., see Bowcock, A.M., 165
Ferreira, A., see Cáceres, A., 233
Figueroa, H.H., see Sauceda, R., 123
Folkesson, R., Monstein, H.-J., Geijer, T., Påhlman, S., Nilsson, K. and Terenius, L., Expression of the proenkephalin gene in human neuroblastoma cell lines, 147
Forloni, G.L., see Guarda, A.S., 223

Fricker, L.D., see Supattapone, S., 173
Friedman, W.J., Dreyfus, C.F., McEwen, B. and Black, I.B., Substance K (NKA) increases tyrosine hydroxylase mRNA in cultured substantia nigra, 203

Gall, C.M., see White, J.D., 21
Garber, S.L. and Makman, M.H., Regulation of tryptophan hydroxylase activity by a cyclic AMP-dependent mechanism in rat striatum, 1
Gardner, P.D., Heinemann, S. and Patrick, J., Transcriptional regulation of nicotinic acetylcholine receptor genes: identification of control elements of a γ-subunit gene, 69
Gariglio, P., see Sauceda, R., 123
Geijer, T., see Folkesson, R., 147
Ginns, E., see Kim, Y.S., 77
Goldstein, M.E., Weiss, S.R., Lazzarini, R.A., Shneidman, P.S., Lees, J.F. and Schlaepfer, W.W., mRNA levels of all three neurofilament protein decline following nerve transection, 287
Goslin, K., see Bowcock, A.M., 165
Greengard, P., see Walaas, S.I., 215
Guarda, A.S., Robinson, M.B., Ory-Lavollée, L., Forloni, G.L., Blakely, R.D. and Coyle, J.T., Quantitation of *N*-acetyl-aspartyl-glutamate in microdissected rat brain nuclei and peripheral tissues: findings with a novel liquid phase radioimmunoassay, 223
Gubits, R.M., Hazelton, J.L. and Simantov, R., Variations in c-fos gene expression during rat brain development, 197
Gutierrez, A.L., see Sauceda, R., 123

Hazelton, J.L., see Gubits, R.M., 197
Heinemann, S., see 69
Heinrich, G., see Allen, J.M., 39
Heinrich, G., see Zheng, M., 133
Herbert, E., see Sivam, S.P., 155
Hoffer, B.., see Läkfors, L., 55
Hoffman, P.N., see Muma, N.A., 115
Hong, J.S., see Sivam, S.P., 155
Hong, J.S., see Stachowiak, M.K., 263
Hong, J.S., see Stachowiak, M.K., 275
Huie, D., see Banner, C., 247
Hwang, J.-J., see Banner, C., 247

Kaczmarek, L., Siedlecki, J.A. and Danysz, W., Proto-oncogene c-fos induction in rat hippocampus, 183
Kage, R., see McGregor, G.P., 207
Kaneko, S., Kato, K., Yamagashi, S., Sugiyama, H. and Nomura, Y., GTP-binding proteins G_i and G_o transplanted onto *Xenopus* oocyte by rat brain messenger RNA, 11
Kaplan, B.B., see Stachowiak, M., 193
Kato, K., see Kaneko, S., 11
Kawai, N., see Saito, M., 83
Kim, Y.S., Thomas, J.W., Tillakaratne, N.J.K., Montpied, P., Suzdak, P.D., Banner, C., Ginns, E., Tobin, A.J. and Paul, S.M., Glutamic acid decarboxylase mRNA in rat brain: regional distribution and effects of intrastriatal kainic acid, 77

Molecular Brain Research, 3 (1988) 298–300
Elsevier

Author Index

Allen, J.M., Martin, J.B. and Heinrich, G., Neuropeptide Y gene expression in PC12 cells and its regulation by nerve growth factor: a model for developmental regulation, 39

Banner, C., Hwang, J.-J., Shapiro, R.A., Wenthold, R.J., Nakatani, Y., Lampel, K.A., Thomas, J.W., Huie, D. and Curthoys, N.P., Isolation of a cDNA for rat brain glutaminase, 247
Banner, C., see Kim, Y.S., 77
Bannon, M.J., Elliott, P.J. and Bunney, E.B., Striatal tachykinin biosynthesis: regulation of mRNA and peptide levels by dopamine agonists and antagonists, 31
Bech-Hansen, N.T., see McLachlan, D.R.C., 255
Bergeron, C., see McLachlan, D.R.C., 255
Biswal, N., Patel, A.G. and Max, S.R., Regulaton of viral and cellular genes in a human neuroblastoma cell line latently infected with herpes simplex virus type 2, 95
Black, I.B., see Friedman, W.J., 203
Blakely, R.D., see Guarda, A.S., 223
Bowcock, A.M., Fahnestock, M., Goslin, K. and Shooter, E.M., The NGF and kallikrein genes of mouse, the African rat *Mastomys natalensis* and man: their distribution and mode of expression in the salivary gland, 165
Brown, I.R., see Sprang, G.K., 89
Brugge, J.S., see Walaas, S.I., 215
Bunney, E.B., see Bannon, M.J., 31
Busciglio, J., see Cáceres, A., 233

Cáceres, A., Busciglio, J., Ferreira, A. and Steward, O., An immunocytochemical and biochemical study of the microtubule-associated protein MAP-2 during post-lesion dendritic remodeling in the central nervous system of adult rats, 233
Calvetta, L., see Sivam, S.P., 155
Carbini, L., see McCaman, R.E., 107
Civelli, O., see Sivam, S.P., 155
Conlon, J.M., see McGregor, G.P., 207
Cooper, B.J., see Roch, J.-M., 61
Coyle, J.T., see Guarda, A.S., 223
Curthoys, N.P., see Banner, C., 247

Danysz, W., see Kaczmarek, L., 183
Davidson, N., see Sutton, F., 187
Davis, L.G., see Manning, R.W., 293
Deguchi, T., see Saito, M., 83
Douglass, J., see Sivam, S.P., 155
Dreyfus, C.F., see Friedman, W.J., 203

Ebendal, T., see Läkfors, L., 55
Elliott, P.J., see Bannon, M.J., 31

Fahnestock, M., see Bowcock, A.M., 165
Ferreira, A., see Cáceres, A., 233
Figueroa, H.H., see Sauceda, R., 123
Folkesson, R., Monstein, H.-J., Geijer, T., Påhlman, S., Nilsson, K. and Terenius, L., Expression of the proenkephalin gene in human neuroblastoma cell lines, 147
Forloni, G.L., see Guarda, A.S., 223

Fricker, L.D., see Supattapone, S., 173
Friedman, W.J., Dreyfus, C.F., McEwen, B. and Black, I.B., Substance K (NKA) increases tyrosine hydroxylase mRNA in cultured substantia nigra, 203

Gall, C.M., see White, J.D., 21
Garber, S.L. and Makman, M.H., Regulation of tryptophan hydroxylase activity by a cyclic AMP-dependent mechanism in rat striatum, 1
Gardner, P.D., Heinemann, S. and Patrick, J., Transcriptional regulation of nicotinic acetylcholine receptor genes: identification of control elements of a γ-subunit gene, 69
Gariglio, P., see Sauceda, R., 123
Geijer, T., see Folkesson, R., 147
Ginns, E., see Kim, Y.S., 77
Goldstein, M.E., Weiss, S.R., Lazzarini, R.A., Shneidman, P.S., Lees, J.F. and Schlaepfer, W.W., mRNA levels of all three neurofilament protein decline following nerve transection, 287
Goslin, K., see Bowcock, A.M., 165
Greengard, P., see Walaas, S.I., 215
Guarda, A.S., Robinson, M.B., Ory-Lavollée, L., Forloni, G.L., Blakely, R.D. and Coyle, J.T., Quantitation of *N*-acetyl-aspartyl-glutamate in microdissected rat brain nuclei and peripheral tissues: findings with a novel liquid phase radioimmunoassay, 223
Gubits, R.M., Hazelton, J.L. and Simantov, R., Variations in c-fos gene expression during rat brain development, 197
Gutierrez, A.L., see Sauceda, R., 123

Hazelton, J.L., see Gubits, R.M., 197
Heinemann, S., see 69
Heinrich, G., see Allen, J.M., 39
Heinrich, G., see Zheng, M., 133
Herbert, E., see Sivam, S.P., 155
Hoffer, B.., see Läkfors, L., 55
Hoffman, P.N., see Muma, N.A., 115
Hong, J.S., see Sivam, S.P., 155
Hong, J.S., see Stachowiak, M.K., 263
Hong, J.S., see Stachowiak, M.K., 275
Huie, D., see Banner, C., 247
Hwang, J.-J., see Banner, C., 247

Kaczmarek, L., Siedlecki, J.A. and Danysz, W., Proto-oncogene c-fos induction in rat hippocampus, 183
Kage, R., see McGregor, G.P., 207
Kaneko, S., Kato, K., Yamagashi, S., Sugiyama, H. and Nomura, Y., GTP-binding proteins G_i and G_o transplanted onto *Xenopus* oocyte by rat brain messenger RNA, 11
Kaplan, B.B., see Stachowiak, M., 193
Kato, K., see Kaneko, S., 11
Kawai, N., see Saito, M., 83
Kim, Y.S., Thomas, J.W., Tillakaratne, N.J.K., Montpied, P., Suzdak, P.D., Banner, C., Ginns, E., Tobin, A.J. and Paul, S.M., Glutamic acid decarboxylase mRNA in rat brain: regional distribution and effects of intrastriatal kainic acid, 77

Koo, E.H., see Muma, N.A., 115

Läkfors, L., Ebendal, T., Whittemore, S.R., Persson, H., Hoffer, B. and Olson, L., Decreased level of nerve growth factor (NGF) and its messenger RNA in the aged rat brain, 55
Lampe, R.A., see Manning, R.W., 293
Lampel, K.A., see Banner, C., 247
Lazzarini, R.A., see Goldstein, M.E., 287
Lee, P.H.K., see Stachowiak, M.K., 263
Lee, P.H.K., see Stachowiak, M.K., 275
Lees, J.F., see Goldstein, M.E., 287
Lester, H.A., see Sutton, F., 187
Li, S., see Sivam, S.P., 155
Lukiw, W.J., see McLachlan, D.R.C., 255
Lustig, A., see Walaas, S.I., 215

Maines, V., see McCaman, R.E., 107
Makman, M.H., see Garber, S.L., 1
Manning, R.W., Reid, C.M., Lampe, R.A. and Davis, L.G., Identification in rodents and other species of an mRNA homologous to the human β-amyloid precursor, 293
Martin, J.B., see Allen, J.M., 39
Matthieu, J.-M., see Roch, J.-M., 61
Max, S.R., see Biswal, N., 95
McCaman, R.E., Carbini, L., Maines, V. and Salvaterra, P.M., Single RNA species injected in *Xenopus* oocyte directs the synthesis of active choline acetyltransferase, 107
McEwen, B., see Friedman, W.J., 203
McGinty, J.F., see Sivam, S.P., 155
McGregor, G.P., Kage, R. and Conlon, J.M., Measurement and partial characterizaton of the C-terminal flanking peptides from bovine preprotachykinins in extracts of brain and gut, 207
McLachlan, D.R.C., Lukiw, W.J., Wong, L., Bergeron, C. and Bech-Hansen, N.T., Selective messenger RNA reduction in Alzheimer's disease, 255
Monstein, H.-J., see Folkesson, R., 147
Montpied, P., see Kim, Y.S., 77
Muma, N.A., Troncoso, J.C., Hoffman, P.N., Koo, E.H. and Price, D.L., Aluminum neurotoxicity: altered expression of cytoskeletal genes, 115

Nakatani, Y., see Banner, C., 247
Nilsson, K., see Folkesson, R., 147
Nomura, Y., see Kaneko, S., 11

Ocadiz, R., see Sauceda, R., 123
Ohsako, S., see Saito, M., 83
Olson, L., see Läkfors, L., 55
Ortega, V., see Sauceda, R., 123
Ory-Lavollée, L., see Guarda, A.S., 223

Påhlman, S., see Folkesson, R., 147
Patel, A.G., see Biswal, N., 95
Patrick, J., see Gardner, P.D., 69
Paul, S.M., see Kim, Y.S., 77
Persson, H., see Läkfors, L., 55
Price, D.L., see Muma, N.A., 115

Ramirez, M., see Roch, J.-M., 61
Reid, C.M., see Manning, R.W., 293
Richardson, P.M., see Riopelle, R.J., 45
Rigual, R.J., see Stachowiak, M.K., 263
Rigual, R.J., see Stachowiak, M.K., 275
Riopelle, R.J., Verge, V.M.K. and Richardson, P.M., Properties of receptors for nerve growth factor in the mature rat nervous system, 45
Robinson, M.B., see Guarda, A.S., 223
Roch, J.-M., Cooper, B.J., Ramirez, M. and Matthieu, J.-M., Expression of only one myelin basic protein allele in mouse is compatible with normal myelination, 61

Saito, M., Ohsako, S., Deguchi, T. and Kawai, N., Glutamate receptors expressed in *Xenopus* oocyte by messenger RNA from invertebrate muscle, 83
Salcedo, M., see Sauceda, R., 123
Salvaterra, P.M., see McCaman, R.E., 107
Sauceda, R., Ocadiz, R., Gutierrez, A.L., Salcedo, M., Ortega, V., Figueroa, H.H. and Gariglio, P., Novel combination of c-myc, N-myc and N-ras oncogene alteraton in brain tumors, 123
Schlaepfer, W.W., see Goldstein, M.E., 287
Schwartz, J.P., Chronic exposure to opiate agonists increases proenkephalin biosynthesis in NG108 cells, 141
Shapiro, R.A., see Banner, C., 247
Shneidman, P.S., see Goldstein, M.E., 287
Shooter, E.M., see Bowcock, A.M., 165
Siedlecki, J.A., see Kaczmarek, L., 183
Simantov, R., see Gubits, R.M., 197
Sivam, S.P., Takeuchi, K., Li, S., Douglas, J., Civelli, O., Calvetta, L., Herbert, E., McGinty, J.F. and Hong, J.S., Lithium increases dynorphin A(1–8) and prodynorphin mRNA levels in the basal ganglia of rats, 155
Snyder, S.H., see Supattapone, S., 173
Sprang, G.K. and Brown, I.R., Selective induction of a heat shock gene in fibre tracts and cerebellar neurons of the rabbit brain detected by in situ hybridization, 89
Stachowiak, M., Stricker, E.M., Zigmond, M.J. and Kaplan, B.B., A cholinergic antagonist blocks cold stress-induced alterations in rat adrenal tyrosine hydroxylase mRNA, 193
Stachowiak, M.K., Lee, P.H.K., Rigual, R.J., Viveros, O.H. and Hong, J.S., Roles of the pituitary-adrenocortical axis in control of the native and cryptic enkephalin levels and proenkephalin mRNA in the sympathoadrenal system of the rat, 263
Stachowiak, N.K., Rigual, R.J., Lee, P.H.K., Viveros, O.H. and Hong, J.S., Regulation of tyrosine hydroxylase and phenylethanolamine *N*-methyltransferase mRNA levels in the sympathoadrenal system by the pituitary-adrenocortical axis, 275
Steward, O., see Cáceres, A., 233
Stricker, E.M., see Stachowiak, M., 193
Strittmatter, S.M., see Supattapone, S., 173
Sugiyama, H., see Kaneko, S., 11
Supattapone, S., Strittmatter, S.M., Fricker, L.D. and Snyder, S.H., Characterization of a neutral, divalent cation-sensitive endopeptidase: a possible role in neuropeptide processing, 173
Sutton, F., Davidson, N. and Lester, H.A., Tetrodotoxin-sensitive voltage-dependent Na currents recorded from *Xenopus* oocytes injected with mammalian cardiac muscle RNA, 187
Suzdak, P.D., see Kim, Y.S., 77

Takeuchi, K., see Sivam, S.P., 155
Terenius, L., see Folkesson, R., 147
Thomas, J.W., see Banner, C., 247

Thomas, J.W., see Kim, Y.S., 77
Tillakaratne, N.J.K., see Kim, Y.S., 77
Tobin, A.J., see Kim, Y.S., 77
Troncoso, J.C., see Muma, N.A., 115

Verge, V.M.K., see Riopelle, R.J., 45
Viveros, O.H., see Stachowiak, M.K., 263
Viveros, O.H., see Stachowiak, M.K., 275

Walaas, S.I., Lustig, A., Greengard, P. and Brugge, J.S., Widespread distribution of the c-*src* gene product in nerve cells and axon terminals in the adult brain, 215
Weiss, S.R., see Goldstein, M.E., 287
Wenthold, R.J., see Banner, C., 247
White, J.D. and Gall, C.M., Differential regulaton of neuropeptide and proto-oncogene mRNA contents in the hippocampus following recurrent seizures, 21
Whittemore, S.R., see Läkfors, L., 55
Wong, L., see McLachlan, D.R.C., 255

Yamagashi, S., see Kaneko, S., 11

Zheng, M. and Heinrich, G., Structural and functional analysis of the promoter region of the nerve growth factor gene, 133
Zigmond, M.J., see Stachowiak, M., 193